



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road

Newport
South Wales
NP10 800
WIPO

21 OCT 2004

PCT

BEST AVAILABLE COPY

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., p.l.c. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed


Dated 11 October 2004

Patents Form 1/77

Patent Act 1977
(Rule 20)



18SEP03 0830090-0029340
P01/7700-000-0321827.8



Request for grant of a patent

18 SEP 2003

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

18 SEP 2003

1. Your reference 101222-1 GB

2. Patent application number
(The Patent Office will fill in this part)

0321827.8

3. Full name, address and postcode of the or of each applicant (underline all surnames)

AstraZeneca UK Limited
15 Stanhope Gate
London
W1K 1LN

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

England

7876467002

4. Title of the invention

CHEMICAL COMPOUNDS

5. Name of your agent (if you have one)

Kevin BILL

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

AstraZeneca
Global Intellectual Property
PO Box 272
Mereside, Alderley Park
Macclesfield,
Cheshire SK10 4GR

Patents ADP number (if you know it)

8179707001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form

Description 8

Claim(s) 2

Abstract

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date 17.09.03

12. Name and daytime telephone number of person to contact in the United Kingdom

Jennifer Bennett - 01625 230148

Warning


After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

CHEMICAL COMPOUND

This invention concerns new polymorphic forms of (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid tris(hydroxymethyl)methylammonium salt (1) (illustrated below), which is useful for the production of a pharmaceutical useful in the treatment of, inter alia, hypercholesterolemia, hyperlipoproteinemia and atherosclerosis.



1

The sodium salt and calcium salt of (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid (hereinafter referred to as compound (2)) were disclosed in European Patent 0521471. This patent also describes a process for the synthesis of the calcium salt, via the sodium salt:

Our International Patent Application WO 00/42024 discloses a crystalline form of the calcium salt of (2), and processes for making it.

Our International Patent Application WO 01/60804 discloses alternative crystalline salts of (2). One of these salts is the tris(hydroxymethyl)methylammonium salt (1). In this application, the process exemplified for formation of tris(hydroxymethyl)methylammonium salt is: acidification of a solution of the methylamine salt of (2) in acetonitrile and water, separation and drying of the organic layer followed by addition of tris(hydroxymethyl)aminomethane at ambient temperature, collection of the crystalline product at ambient temperature and then drying of the crystals at 30°C under vacuum. This process produces needle shaped crystals of a single polymorph of the salt (1) with an X-ray powder diffraction pattern with peaks at 2-theta = 7.9, 8.5, 10.2, 16.7, 18.4, 19.3, 19.8, 20.2, 21.5 and 24.9°.

We have discovered two further polymorphic crystalline forms of the tris(hydroxymethyl)methylammonium salt (1) herein called Forms 2 and 3. Such polymorphic forms may have different solubilities and/or stabilities and/or bioavailabilities

and/or different impurity profiles (minor impurities which arise for example because of the process of manufacture and/or isolation) and/or crystal forms which are easier to handle, micronise and/or form into tablets.

According to one aspect of the invention is provided a crystalline
5 tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern with peaks at 2-theta = 3.2, 6.3, 9.5 and 11.0. This crystalline form is hereinafter referred to as Form 2.

According to another aspect of the invention is provided a crystalline
10 tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern with peaks at 2-theta = 3.2, 6.3, 9.5, 11.0, 12.0, 12.4, 13.9 and 21.5.

According to another aspect of the invention is provided a crystalline
15 tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern with peaks at 2-theta = 3.2, 6.3, 9.5, 11.0, 12.0, 12.4, 13.9, 15.8, 21.5, 22.7, 23.6 and 24.9.

According to another aspect of the invention is provided a crystalline
20 tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern substantially as shown in Figure 1.

It will be appreciated that the 2-theta values listed in the aspects of the invention hereinbefore for Form 2, and hereinafter for Form 3, are chosen because they most clearly
25 differentiate one Form from another, although they do not necessarily represent the most intense peaks.

The Form 2 polymorphic salt of this aspect of the invention may be produced by the following process: a sample of amorphous tris(hydroxymethyl)methylammonium salt (1) (which may be prepared by freeze drying an aqueous solution of the salt (1)) is slurried in a
30 suitable organic solvent at a temperature below ambient temperature, the resultant mixture is filtered and the resulting product is dried as necessary.

Suitable organic solvents may be determined experimentally by the skilled person. Conveniently, the organic solvent is acetonitrile, ethyl acetate or MTBE (methylt-butylether).

Conveniently the mixture is slurried for an extended period, for example for 24 hours. Conveniently, the mixture is slurried at a temperature below ambient temperature which is for example, between about 0°C and 10°C, such as between about 0°C and 5°C, and preferably at about 0°C.

5 The product is conveniently dried by prolonged filtration under vacuum, the use of temperatures above ambient temperature preferably being avoided in order to avoid any risk of conversion of polymorphic form.

According to a further aspect of the invention is provided a crystalline tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-
10 [methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern with peaks at 2-theta = 6.9 and 13.1. This crystalline form is hereinafter referred to as Form 3.

According to a further aspect of the invention is provided a crystalline tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-
15 [methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern with peaks at 2-theta = 6.9, 13.1, 14.9 and 20.6.

According to a further aspect of the invention is provided a crystalline tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-
20 [methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern with peaks at 2-theta = 6.9, 8.5, 9.0, 13.1, 14.9, 17.2, 18.2, 18.6, 19.0, 19.4, 20.6 and 25.4.

According to another aspect of the invention is provided a crystalline tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-
25 [methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern substantially as shown in Figure 2.

The Form 3 polymorphic salt of this aspect of the invention may be produced by the following process: a sample of amorphous tris(hydroxymethyl)methylammonium salt (1) (which may be prepared by freeze drying an aqueous solution of the salt (1)) is slurried in isopropanol at a temperature below ambient temperature, the resultant mixture is filtered and
30 the resulting product is dried.

Conveniently the mixture is slurried for an extended period, for example for 24 hours. Conveniently, the mixture is slurried at a temperature below ambient temperature which is,

for example, between about 0°C and 10°C, such as between about 0°C and 5°C, and preferably at about 0°C.

The product is conveniently dried by prolonged filtration under vacuum, the use of temperatures above ambient temperature preferably being avoided in order to avoid any risk of conversion of polymorphic form.

Thermal Gravimetric Analysis of samples of Form 3 indicates that the polymorphic form is solvated, which arises from the method of manufacture and will be water and/or isopropanol.

The X-ray powder diffraction spectra were determined by mounting a sample of the crystalline form on Siemens single silicon crystal (SSC) wafer mounts and spreading out the sample into a thin layer with the aid of a microscope slide. The sample was spun at 30 revolutions per minute (to improve counting statistics) and irradiated with X-rays generated by a copper long-fine focus tube operated at 40kV and 40mA with a wavelength of 1.5406 angstroms. The collimated x-ray source was passed through an automatic variable divergence slit set at V20 (20mm path length) and the reflected radiation directed through a 2mm antiscatter slit and a 0.2mm detector slit. The sample was exposed for 4 seconds per 0.02 degree 2-theta increment (continuous scan mode) over the range 2 degrees to 40 degrees 2-theta in theta-theta mode. The running time was 2 hours 6 minutes and 40 seconds. The instrument was equipped with a scintillation counter as detector. Control and data capture was by means of a DECpc LPv 433sx personal computer running with Diffrac AT (Socabim) software.

It will be understood that the 2-theta values of an X-ray powder diffraction pattern may vary slightly from one machine to another or from one sample to another, and so the values quoted are not to be construed as absolute. It will also be understood that the relative intensities of peaks may vary according to the orientation of the sample under test so that the intensities in the XRD traces included herein are illustrative and not intended to be used for absolute comparison.

Forms 2 and 3 obtained according to the present invention are substantially free from other crystal and non-crystal forms of tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid. The term "substantially free from other crystal and non-crystal forms" shall be understood to mean that the desired crystal form (Form 2 or Form 3) contains less than 50%, preferably less than 10%, more preferably less than 5% of any other forms of

the tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid.

The utility of the compounds of the invention may be demonstrated by standard tests and clinical studies, including those described in EPA 521471.

5 According to a further feature of the invention is a method of treating a disease condition wherein inhibition of HMG CoA reductase is beneficial which comprises administering to a warm-blooded mammal an effective amount of Form 2 or Form 3. The invention also relates to the use of Form 2 or Form 3 in the manufacture of a medicament for use in a disease condition.

10 The compound of the invention may be administered to a warm-blooded animal, particularly a human, in need thereof for treatment of a disease in which HMG CoA reductase is implicated, in the form of a conventional pharmaceutical composition. Therefore in another aspect of the invention, there is provided a pharmaceutical composition comprising Form 2 or Form 3 in admixture with a pharmaceutically acceptable carrier.

15 Such compositions may be administered in standard manner for the disease condition that it is desired to treat, for example by oral, topical, parenteral, buccal, nasal, vaginal or rectal administration or by inhalation. For these purposes Form 2 or Form 3 may be formulated by means known in the art into the form of, for example, tablets, capsules, aqueous or oily solutions, suspensions, emulsions, creams, ointments, gels, nasal sprays, 20 suppositories, finely divided powders or aerosols for inhalation, and for parenteral use (including intravenous, intramuscular or infusion) sterile aqueous or oily solution or suspensions or sterile emulsions. A preferred route of administration is oral. Form 2 or Form 3 will be administered to humans at a daily dose in, for example, the ranges set out in EPA 521471. The daily doses may be given in divided doses as necessary, the precise amount of 25 the Form received and the route of administration depending on the weight, age and sex of the patient being treated and on the particular disease condition being treated according to principles known in the art.

 According to a further feature of the invention, there is provided a process for the manufacture of a pharmaceutical composition containing Form 2 or Form 3 as active 30 ingredient, which comprises admixing Form 2 or Form 3 together with a pharmaceutically acceptable carrier.

The invention is further illustrated, but not limited by the following examples.

Example 1

Amorphous tris(hydroxymethyl)methylammonium salt (1) (1 g), prepared by freeze-drying an aqueous solution of the salt, was added to acetonitrile (10 ml) at 0°C and stirred at 0°C for 24 h. The slurry was filtered under vacuum to dryness to yield

5 tris(hydroxymethyl)methylammonium salt (1) Form 2.

Example 2

Amorphous tris(hydroxymethyl)methylammonium salt (1) (1 g), prepared by freeze-drying an aqueous solution of the salt, was added to ethyl acetate (10 ml) at 0°C and stirred at 0°C for

10 24 h. The slurry was filtered under vacuum to dryness to yield
tris(hydroxymethyl)methylammonium salt (1) Form 2.

Example 3


Amorphous tris(hydroxymethyl)methylammonium salt (1) (1 g), prepared by freeze-drying an aqueous solution of the salt, was added to MTBE (10 ml) at 0°C and stirred at 0°C for 24 h.

15 The slurry was filtered under vacuum to dryness to yield
tris(hydroxymethyl)methylammonium salt (1) Form 2.

Example 4


20 Amorphous tris(hydroxymethyl)methylammonium salt (1) (1 g), prepared by freeze-drying an aqueous solution of the salt, was added to isopropyl alcohol (10 ml) at 0°C and stirred at 0°C for 24 h. The slurry was filtered under vacuum to dryness to yield
tris(hydroxymethyl)methylammonium salt (1) Form 3.

Figure 1. Tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid Form 2



2-theta	d-spacing	Relative Intensity
3.2	27.8	29
6.3	14.0	18
9.5	9.3	54
11.0	8.0	99
12.0	7.4	14
12.4	7.2	31
13.9	6.4	51
15.8	5.6	22
21.5	4.1	100
22.7	3.9	25
23.6	3.8	19
24.9	3.6	16


Figure 2 – Tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid Form 3



2-theta	d-spacing	Relative Intensity
6.9	12.8	100
8.5	10.5	41
9.0	9.9	12
13.1	6.7	13
14.9	6.0	24
17.2	5.1	58
18.2	4.9	58
18.6	4.8	39
19.0	4.7	38
19.4	4.6	30
20.6	4.3	63
25.4	3.5	30

Claims

1. A crystalline form of the compound tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid of the formula (I) having an X-ray powder diffraction pattern with specific peaks at 2-theta = 3.2, 6.3, 9.5 and 11.0.



(I)

2. A crystalline form as claimed in Claim 1 having an X-ray powder diffraction pattern with specific peaks at 2-theta = 3.2, 6.3, 9.5, 11.0, 12.0, 12.4, 13.9 and 21.5.
3. A crystalline form as claimed in Claim 1 having an X-ray powder diffraction pattern with specific peaks at 2-theta = 3.2, 6.3, 9.5, 11.0, 12.0, 12.4, 13.9, 15.8, 21.5, 22.7, 23.6 and 24.9.
4. A crystalline form of the compound tris(hydroxymethyl)methylammonium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid having an X-ray powder diffraction pattern with specific peaks at 2-theta = 6.9 and 13.1.
5. A crystalline form as claimed in Claim 4 having an X-ray powder diffraction pattern with specific peaks at 2-theta = 6.9, 13.1, 14.9 and 20.6.
6. A crystalline form as claimed in Claim 4 having an X-ray powder diffraction pattern with specific peaks at 2-theta = 6.9, 8.5, 9.0, 13.1, 14.9, 17.2, 18.2, 18.6, 19.0, 19.4, 20.6 and 25.4.

5. A pharmaceutical composition comprising a crystalline form as claimed in any one of the preceding claims, together with a pharmaceutically acceptable carrier.

6. A process for the manufacture of a pharmaceutical composition as claimed in claim 5 which comprises admixing a crystalline form as claimed in Claim 1 or Claim 4 together with a pharmaceutically acceptable carrier.

7. The use of a crystalline form as claimed in Claim 1 or Claim 4 in the manufacture of a medicament.

8. A method of treating a disease condition wherein inhibition of HMG CoA reductase is beneficial which comprises administering to a warm-blooded mammal an effective amount of a crystalline form as claimed in Claim 1 or Claim 4.

9. A process for the manufacture of a crystalline form as claimed in Claim 1 or Claim 4 which comprises forming crystals by:

a) slurring a sample of amorphous tris(hydroxymethyl)methylammonium salt (1) in an organic solvent at a temperature below ambient temperature;

b) filtration of the resultant mixture; and

c) drying of the resultant product as necessary.

10. A process as claimed in Claim 9 for the manufacture of Form 2 wherein the organic solvent is acetonitrile, ethyl acetate or MTBE (methylt-butylether).

11. A process for the manufacture of a crystalline form as claimed in Claim 9 for the manufacture of Form 3 which comprises forming crystals by:

a) slurring a sample of amorphous tris(hydroxymethyl)methylammonium salt (1) in isopropanol at a temperature below ambient temperature;

b) filtration of the resultant mixture; and

c) drying of the resultant product as necessary.

12. A process as claimed in any one of Claims 9 to 11 wherein the temperature is about 0°C.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.